

Appendix M-II-B-1-a. What is the structure of the cloned DNA that will be used?

Appendix M-II-B-1-a-(1). Describe the gene (genomic or cDNA), the bacterial plasmid or phage vector, and the delivery vector (if any). Provide complete nucleotide sequence analysis or a detailed restriction enzyme map of the total construct.

Appendix M-II-B-1-a-(2). What regulatory elements does the construct contain (e.g., promoters, enhancers, polyadenylation sites, replication origins, etc.)? From what source are these elements derived? Summarize what is currently known about the regulatory character of each element.

Appendix M-II-B-1-a-(3). Describe the steps used to derive the DNA construct.

Appendix M-II-B-1-b. What is the structure of the material that will be administered to the patient?

Appendix M-II-B-1-b-(1). Describe the preparation, structure, and composition of the materials that will be given to the patient or used to treat the patient's cells: (i) If DNA, what is the purity (both in terms of being a single DNA species and in terms of other contaminants)? What tests have been used and what is the sensitivity of the tests? (ii) If a virus, how is it prepared from the DNA construct? In what cell is the virus grown (any special features)? What medium and serum are used? How is the virus purified? What is its structure and purity? What steps are being taken (and assays used with their sensitivity) to detect and eliminate any contaminating materials (for example, VL30 RNA, other nucleic acids, or proteins) or contaminating viruses (both replication-competent or replication-defective) or other organisms in the cells or serum used for preparation of the virus stock including any contaminants that may have biological effects? (iii) If co-cultivation is employed, what kinds of cells are being used for co-cultivation? What steps are being taken (and assays used with their sensitivity) to detect and eliminate any contaminating materials? Specifically, what tests are being conducted to assess the material to be returned to the patient for the presence of live or killed donor cells or other non-vector materials (for example, VL30 sequences) originating from those cells? (iv) If methods other than those covered by Appendices M-II-B-1 through M-II-B-3 are used to introduce new genetic information into target cells, what steps are being taken to detect and eliminate any contaminating materials? What are possible sources of contamination?

What is the sensitivity of tests used to monitor contamination?

Appendix M-II-B-1-b-(2). Describe any other material to be used in preparation of the material to be administered to the patient. For example, if a viral vector is proposed, what is the nature of the helper virus or cell line? If carrier particles are to be used, what is the nature of these?

Appendix M-II-B-2. Preclinical Studies, Including Risk-Assessment Studies

Provide results that demonstrate the safety, efficacy, and feasibility of the proposed procedures using animal and/or cell culture model systems, and explain why the model(s) chosen is/are most appropriate.

Appendix M-II-B-2-a. Delivery System

Appendix M-II-B-2-a-(1). What cells are the intended target cells of recombinant DNA? What target cells are to be treated *ex vivo* and returned to the patient, how will the cells be characterized before and after treatment? What is the theoretical and practical basis for assuming that only the target cells will incorporate the DNA?

Appendix M-II-B-2-a-(2). Is the delivery system efficient? What percentage of the target cells contain the added DNA?

Appendix M-II-B-2-a-(3). How is the structure of the added DNA sequences monitored and what is the sensitivity of the analysis? Is the added DNA extrachromosomal or integrated? Is the added DNA unrearranged?

Appendix M-II-B-2-a-(4). How many copies are present per cell? How stable is the added DNA both in terms of its continued presence and its structural stability?

Appendix M-II-B-2-b. Gene Transfer and Expression

Appendix M-II-B-2-b-(1). What animal and cultured cell models were used in laboratory studies to assess the *in vivo* and *in vitro* efficacy of the gene transfer system? In what ways are these models similar to and different from the proposed human treatment?

Appendix M-II-B-2-b-(2). What is the minimal level of gene transfer and/or expression that is estimated to be necessary for the gene transfer protocol to be successful in humans? How was this level determined?

Appendix M-II-B-2-b-(3). Explain in detail all results from animal and cultured cell model experiments which assess the effectiveness of the delivery system in achieving the minimally

required level of gene transfer and expression.

Appendix M-II-B-2-b-(4). To what extent is expression only from the desired gene (and not from the surrounding DNA)? To what extent does the insertion modify the expression of other genes?

Appendix M-II-B-2-b-(5). In what percentage of cells does expression from the added DNA occur? Is the product biologically active? What percentage of normal activity results from the inserted gene?

Appendix M-II-B-2-b-(6). Is the gene expressed in cells other than the target cells? If so, to what extent?

Appendix M-II-B-2-c. Retrovirus Delivery Systems

Appendix M-II-B-2-c-(1). What cell types have been infected with the retroviral vector preparation? Which cells, if any, produce infectious particles?

Appendix M-II-B-2-c-(2). How stable are the retroviral vector and the resulting provirus against loss, rearrangement, recombination, or mutation? What information is available on how much rearrangement or recombination with endogenous or other viral sequences is likely to occur in the patient's cells? What steps have been taken in designing the vector to minimize instability or variation? What laboratory studies have been performed to check for stability, and what is the sensitivity of the analyses?

Appendix M-II-B-2-c-(3). What laboratory evidence is available concerning potential harmful effects of the transfer (e.g., development of neoplasia, harmful mutations, regeneration of infectious particles, or immune responses)? What steps will be taken in designing the vector to minimize pathogenicity? What laboratory studies have been performed to check for pathogenicity, and what is the sensitivity of the analyses?

Appendix M-II-B-2-c-(4). Is there evidence from animal studies that vector DNA has entered untreated cells, particularly germ-line cells? What is the sensitivity of these analyses?

Appendix M-II-B-2-c-(5). Has a protocol similar to the one proposed for a clinical trial being conducted in non-human primates and/or other animals? What were the results? Specifically, is there any evidence that the retroviral vector has recombined with any endogenous or other viral sequences in the animals?