

TABLE 1.—1995 SPECIFICATIONS OF ACCEPTABLE BIOLOGICAL CATCH (ABC), HARVEST GUIDELINES, AND LIMITED-ENTRY AND OPEN-ACCESS ALLOCATIONS, BY INTERNATIONAL NORTH PACIFIC FISHERIES COMMISSION (INPFC) SUBAREAS

| Species | Acceptable Biological Catch (ABC) (x1,000 mt) | | | | | | Total ABC | Harvest Guideline (x1,000 mt) | Allocations (x1,000 mt) | | | |
|--------------------------------|---|------------------|------------------|------------------|------------------|---------|-------------------|-------------------------------|-------------------------|--------------|--------------|-------------|
| | Van-couver ^a | Columbia | Eureka | Monterey | Concept-ion | 1000 mt | | | Percent | 1000 mt | Percent | |
| | | | | | | | | | | | | Open access |
| Roundfish: | | | | | | | | | | | | |
| Lingcod ^b | 1.3 | | 0.3 | 0.7 | 0.1 | | 2.4 | 2.4 | 1.21 | 80.9 | 0.29 | 19.1 |
| Pacific cod | 3.2 | | (^c) | (^c) | (^c) | | | | | | | |
| Pacific whiting ^d | | | 223.0 | | | | 178.4 | 178.4 | | | | |
| Sablefish ^{e,f} | | | 8.7 | | 0.425 | | 9.1 | 7.1 | 5.90 | 93.4 | 0.42 | 6.6 |
| Jack mackerel ^g | | | 52.6 | | | | 52.6 | 52.6 | | | | |
| Rockfish: | | | | | | | | | | | | |
| POP ^h | 0.0 | 0.0 | (^c) | (^c) | (^c) | | 1.3 | 1.3 | | | | |
| Shorthelly | | | 23.5 | | | | 23.5 | 23.5 | | | | |
| Widow ⁱ | | | 7.7 | | | | 7.7 | 6.5 | 6.26 | 96.3 | 0.24 | 3.7 |
| Thornyheads: | | | | | | | | | | | | |
| Shortspine ^{e,j} | | | 8.0 | | | | 8.0 | | | | | |
| Longspine ^{e,j} | | | 1.0 | | | | 1.0 | 1.5 | | | | |
| Sebastes complex: ^k | | | 7.0 | | | | 7.0 | 6.0 | | | | |
| | 11.9 | | | 13.2 | | | 11.9 N, 13.2 S | 11.8 N, 13.2 S | 10.67 8.76 | 90.4 67.4 | 1.13 4.24 | 9.6 32.6 |
| Bocaccio ^l | (^c) | (^c) | | 1.7 | | | 1.7 | 1.7 | 1.01 | 67.4 | 0.49 | 32.6 |
| Canary ^m | 1.0 | | 0.25 | (^c) | (^c) | | 1.25 | 0.85 | 0.78 | 91.2 | 0.07 | 8.8 |
| Chilipepper | (^c) | (^c) | | 4.0 | | | 4.0 | | | | | |
| Yellowtail ⁿ | 1.19 | 2.97 | 2.58 | (^c) | (^c) | | 6.74 | 4.16 N 2.58 S | 3.76 2.33 | 90.4 90.4 | 0.40 0.25 | 9.6 9.6 |
| Remaining rockfish | 0.8 | 3.7 | | 7.0 | | | 11.5 | | | | | |
| Flatfish: | | | | | | | | | | | | |
| Dover sole ^{e,o} | 2.4 | 3.0 | 2.9 | 5.0 | 1.0 | | 14.3 | 13.6 WOC, 2.85 Col | | | | |
| English sole | 2.0 | | | 1.1 | | | 3.1 | | | | | |
| Petrals sole | 1.2 | | 0.5 | 0.8 | 0.2 | | 2.7 | | | | | |
| Arrowtooth Flounder | | | 5.8 | | | | 5.8 | | | | | |
| Other flatfish | 0.7 | 3.0 | 1.7 | 1.8 | 0.5 | | 7.7 | | | | | |
| Other fish ^p | 2.5 | 7.0 | 1.2 | 2.0 | 2.0 | | 14.7 | | | | | |

^a U.S. Vancouver only, except for Pacific whiting.